

SEMINAR

Grupo de Análise Funcional e Aplicações Functional Analysis and Applications Group

Exponentials rarely maximize Fourier extension estimates to cones

Giuseppe Negro

CAMGSD, Instituto Superior Técnico de Lisboa

Abstract

We consider the general Fourier extension estimate to the d -dim. cone in \mathbb{R}^{1+d} :

$$\|\widehat{f\mu}\|_{L^q(\mathbb{R}^{1+d})} \leq C_{d,p} \|f\|_{L^p(d\mu)}.$$

Firstly, we prove that it admits *maximizers*, i.e. functions f that attain the best possible constant $C_{d,p}$. For $p = 2$, this estimate coincides with the *Strichartz estimate* for the wave equation; in this case, Foschi found that $f(\tau, \xi) = \exp(-\tau)$ is a maximizer for $d \in \{2, 3\}$. We further prove that such exponential functions have a chance of being maximizers for general d if and only if $p = 2$. This parallels the analogous result of Christ and Quilodrán that gaussian functions rarely maximize the Fourier extension estimate to the paraboloid. Our proof is entirely different, and it is based on the conformal compactification of \mathbb{R}^{1+d} given by the Penrose transform.

Room 11.2.25
February 23, 2023 - 14:00

This seminar is supported in part by the Portuguese Foundation for Science and Technology (FCT - Fundação para a Ciência e a Tecnologia), through CIDMA - Center for Research and Development in Mathematics and Applications, within project UIDB/04106/2020.